

Adiabatic Geometrical Phase for Scalar Fields in a Curved Spacetime

Ali Mostafazadeh*

Theoretical Physics Institute, University of Alberta,
Edmonton, Alberta T6G 2J1, Canada,

and

Department of Mathematics, College of Arts and Sciences,
Koç University, Istinye, 80860 Istanbul, Turkey.

June 1997

Abstract

A convenient framework is developed to generalize Berry's investigation of the adiabatic geometrical phase for a classical relativistic charged scalar field in a curved background spacetime which is minimally coupled to electromagnetism and an arbitrary (non-electromagnetic) scalar potential. It involves a two-component formulation of the corresponding Klein-Gordon equation. A precise definition of the adiabatic approximation is offered and conditions of its validity are discussed. It is shown that the adiabatic geometric phase can be computed without making a particular choice for an inner product on the space of solutions of the field equations. What is needed is just an inner product on the Hilbert space of the square integrable functions defined on the spatial hypersurfaces. The two-component formalism is applied in the investigation of the adiabatic geometric phases for several specific examples, namely, a rotating magnetic field in Minkowski space, a rotating cosmic string, and an arbitrary spatially homogeneous cosmological background. It is shown that the two-component formalism reproduces the known results for the first two examples. It also leads to several interesting results for the case of spatially homogeneous cosmological models. In particular, it is shown that the adiabatic geometric phase angles vanish for Bianchi type I models. The situation is completely different for Bianchi type IX models where a variety of nontrivial non-Abelian adiabatic geometrical phases can occur. The analogy between the adiabatic geometric phases induced by the Bianchi type IX backgrounds and those associated with the well-known time-dependent nuclear quadrupole Hamiltonians is also pointed out.

Number of pages = 39 + one figure

*E-mail: alimos@phys.ualberta.ca

- **Proposed Running head:** Geometric phase of scalar fields
- **Author:** Ali Mostafazadeh
- **Contact Information:**

- **Before June 27, 1997:**

Address: 412 Physics Lab, University of Alberta, Edmonton, AB, T6G 2J1, Canada

Phone: (403) 434-5877, (403) 492-3960

Fax: (403) 492-0714

- **After June 27, 1997:**

Address: Department of Mathematics, College of Arts and Sciences, Koç University,
Çayir Cad. no: 5, Istinye, 80860 Istanbul, Turkey

Fax: (90-212) 229-0680

- **E-mail:** alimos@phys.ualberta.ca

1 Introduction

Following the pioneering works of Berry [1], Wilczek and Zee [2] and Aharonov and Anandan [3] on geometric phases in non-relativistic quantum mechanics, Garrison and Chiao [4] showed that the geometric phases may be defined for any classical field theory provided that some gauge symmetries were present. The role of the gauge symmetry was to provide a conserved charge which in turn defined an inner product on the function space of the classical fields. Anandan [5], commented on the latter article, indicating that the condition of gauge symmetry may in general be relaxed or practically replaced with the condition of the existence of an inner product on the space of fields. A common assumption of both Refs. [4] and [5] was that the field equations involved only first time derivatives of the fields which is really not a restriction.

Clearly the simplest classical field theories of interest are the Klein-Gordon fields on the ordinary Minkowski spacetime. The phenomenon of the geometric phase for charged Klein-Gordon fields minimally coupled to a time-dependent electromagnetic field has already been studied by Anandan and Mazur [6]. The main strategy of [6] is to decompose the vector space of the fields into three subspaces which are spanned respectively by the positive, zero, and negative frequency (energy) solutions and to note that on the positive and negative frequency subspaces, where the Klein-Gordon inner product is positive, respectively, negative definite, the Klein-Gordon equation may be written as a pair of equations which are linear in the time-derivative of the field. More recently, a similar approach has been pursued to study the dynamics of Klein-Gordon fields in a periodic Friedmann-Robertson-Walker background by Droz-Vincent [7].

The idea of investigating the manifestations of the geometric phase in the context of gravitation and cosmology is in fact not quite new. The first developments in this direction, to best of my knowledge, goes back to the work of Brout and Venturi [8] which was inspired by the earlier results of Banks [9] and Brout [10] on the use of Born-Oppenheimer approximation in semiclassical treatment of the Wheeler-DeWitt equation, and the application of Berry's phase in improving the Born-Oppenheimer approximation in molecular physics [11]. Subsequent work which followed essentially the same idea is that of Venturi [12, 13], Casadio and Venturi [14]

and Datta [15]. There is also the contributions of Cai and Papini [16] which are based on the four-space-formalism of the relativistic quantum mechanics. More recently Corichi and Pierri [17] considered Klein-Gordon fields in a class of stationary spacetimes and in particular investigated the induced topological Aharonov-Bohm type phases due to a rotating cosmic string. The analogy between the topological phase due to a rotating cosmic string and the Aharonov-Bohm phase had previously been pointed out by de Sousa Gerbert and Jackiw [18].

The purpose of this article is to study the geometric phases associated with a charged Klein-Gordon field Φ in an arbitrary globally hyperbolic spacetime (M, g) which is minimally coupled to an electromagnetic potential A , as well as an arbitrary scalar potential V . The latter may, for instance, be identified with the appropriate multiple of the Ricci scalar curvature which renders the theory conformally invariant. The problem of investigation of the dynamics of such a field theory has a long history in the context of developing quantum field theories in a curved background [19, 20, 21]. However, I shall not be concerned with subtleties associated with the full second quantized theory. Instead, the Klein-Gordon field will be viewed and treated as a classical (first quantized) field.

In section 2, the two-component form of the field equation is derived. This allows for a simple application of the adiabatic theorem which following Berry's original approach [1] yields the definition of the adiabatically cyclic states and the associated dynamical and geometric phases for the theory. This is described in section 3 and applied to the problems of a rotating magnetic field in Minkowski space in section 4, a rotating cosmic string in section 5, and spatially homogeneous (Bianchi) cosmological backgrounds in section 6. The latter section also includes a detailed analysis of Bianchi type I and IX cases. Particularly interesting is the analogy between the case of a Bianchi type IX background and the quadrupole Hamiltonians of the molecular and nuclear physics.

2 Two-Component Formalism

Consider complex scalar fields Φ defined on a globally hyperbolic spacetime $(M, g) = (\mathbb{R} \times \Sigma, g)$ satisfying

$$\left[g^{\mu\nu} (\nabla_\mu + ieA_\mu) (\nabla_\nu + ieA_\nu) + V - \mu^2 \right] \Phi = 0 , \quad (1)$$

where $g^{\mu\nu}$ are components of the inverse of the metric g , ∇_μ is the covariant derivative along $\partial/\partial x^\mu$ defined by the Levi Civita connection, A_μ are components of the electromagnetic potential, V is an arbitrary scalar potential, e is the electric charge, and μ is the mass. Throughout this article the signature of the metric g is chosen to be $(-, +, +, +)$, letters from the beginning and the middle of the Greek alphabet are associated with an arbitrary local basis and a local coordinate basis of the tangent spaces (bundle) of the spacetime manifold, respectively. The letters from the beginning and the middle of the Latin alphabet label the corresponding spatial components. They take 1, 2 and 3.

Denoting a time derivative by a dot, one can express Eq. (1) in the form:

$$\ddot{\Phi} + \hat{D}_1 \dot{\Phi} + \hat{D}_2 \Phi = 0 , \quad (2)$$

where

$$\hat{D}_1 := \frac{2}{g^{00}} \left[g^{0i} \partial_i + ie g^{0\mu} A_\mu - \frac{1}{2} g^{\mu\nu} \Gamma_{\mu\nu}^0 \right] , \quad (3)$$

$$\begin{aligned} \hat{D}_2 := & \frac{2}{g^{00}} \left[\frac{1}{2} g^{ij} \partial_i \partial_j + (ie g^{\mu i} A_\mu - \frac{1}{2} g^{\mu\nu} \Gamma_{\mu\nu}^i) \partial_i + \right. \\ & \left. \frac{1}{2} g^{\mu\nu} (ie \nabla_\mu A_\nu - e^2 A_\mu A_\nu) + \frac{1}{2} (V - \mu^2) \right] . \end{aligned} \quad (4)$$

A two-component representation of the field equation (2) is

$$i \dot{\Psi}^{(q)} = \hat{H}^{(q)} \Psi^{(q)} , \quad (5)$$

where

$$\Psi^{(q)} := \begin{pmatrix} u^{(q)} \\ v^{(q)} \end{pmatrix} , \quad (6)$$

$$u^{(q)} := \frac{1}{\sqrt{2}} (\Phi + q \dot{\Phi}) , \quad v^{(q)} := \frac{1}{\sqrt{2}} (\Phi - q \dot{\Phi}) , \quad (7)$$

$$\hat{H}^{(q)} := \frac{i}{2} \begin{pmatrix} \frac{\dot{q}}{q} + \frac{1}{q} - \hat{D}_1 - q \hat{D}_2 & -\frac{\dot{q}}{q} - \frac{1}{q} + \hat{D}_1 - q \hat{D}_2 \\ -\frac{\dot{q}}{q} + \frac{1}{q} + \hat{D}_1 + q \hat{D}_2 & \frac{\dot{q}}{q} - \frac{1}{q} - \hat{D}_1 + q \hat{D}_2 \end{pmatrix} , \quad (8)$$

and q is an arbitrary, possibly time-dependent, non-zero complex parameter. The set $\mathbb{C} - \{0\}$ of q 's defines a group of transformations

$$\Psi^{(q)} \rightarrow \Psi^{(q')} =: g(q', q) \Psi^{(q)} \quad (9)$$

which is isomorphic to $GL(1, \mathbb{C})$. The group elements are given by

$$g(q', q) = g(\gamma) = \begin{pmatrix} \frac{1+\gamma}{2} & \frac{1-\gamma}{2} \\ \frac{1-\gamma}{2} & \frac{1+\gamma}{2} \end{pmatrix},$$

where $\gamma := q'/q$. Under the transformation (9), the Hamiltonian transforms according to

$$\hat{H}^{(q)} \rightarrow \hat{H}^{(q')} = g(\gamma) \hat{H}^{(q)} g^{-1}(\gamma) + i\dot{g}(\gamma) g^{-1}(\gamma),$$

and the Schrödinger equation (5) preserves its form. The underlying $GL(1, \mathbb{C})$ symmetry which characterizes the arbitrariness of q does not have any physical significance. It is, however, useful for computational purposes. A concrete example is given in section 6.

The advantage of the two-component form of the field equation is that it enables one to proceed in a manner analogous with the well-known non-relativistic quantum mechanical case. Indeed Eq. (5) with a fixed choice of q is a Schrödinger equation associated with an explicitly time-dependent Hamiltonian $\hat{H}^{(q)}$. The two-component fields $\Psi^{(q)}$ belong to the vector space $\mathcal{H}_t \oplus \mathcal{H}_t$ where $\mathcal{H}_t$ is the Hilbert space completion (with respect to an appropriate inner product) of compactly supported complex-valued functions on the spatial hypersurface Σ_t associated with a specific ADM decomposition of the spacetime [22].

Usually in the two-component approach to the Klein-Gordon field theory in Minkowski spacetime, one chooses an inner product on $\mathcal{H}_t \oplus \mathcal{H}_t$ in such a way as to make the corresponding Hamiltonian self-adjoint [23, 24]. A Hermitian inner product $(\ , \)$ on $\mathcal{H}_t \oplus \mathcal{H}_t$ may be defined by a Hermitian inner product $\langle \ | \ \rangle$ on $\mathcal{H}_t$ and a possibly time-dependent complex Hermitian 2×2 matrix $h = (h_{rs})$:

$$(\Psi_1, \Psi_2) := (\langle u_1 |, \langle v_1 |) \begin{pmatrix} h_{11} & h_{12} \\ h_{12}^* & h_{22} \end{pmatrix} \begin{pmatrix} |u_2\rangle \\ |v_2\rangle \end{pmatrix}, \quad (10)$$

where u_r and v_r are components of Ψ_r , and h_{11} and h_{22} are real. The usual choice for h , in the Minkowski case, is $h_{11} = -h_{22} = 1$, $h_{12} = 0$, [23, 24]. This choice leads to

$$(\Psi_1, \Psi_2) = \langle u_1 | u_2 \rangle - \langle v_1 | v_2 \rangle. \quad (11)$$

It is not difficult to check that in the general case this choice does not guarantee the self-adjointness of the Hamiltonian unless some severe conditions are imposed on q and the operators $\hat{D}_1$ and $\hat{D}_2$, namely, that q must be imaginary, $\hat{D}_2$ must be self-adjoint with respect to the inner product $\langle | \rangle$ on $\mathcal{H}_t$, and $\hat{D}_1 = \dot{q}/q$. The latter condition is especially restrictive as q can only depend on time and being a free (non-dynamical) parameter, may be set to a constant in which case $\hat{D}_1$ must vanish. In general, these conditions are not fulfilled. Nevertheless, the inner product (11) has an appealing property which is described next.

Consider the eigenvalue problem for $H^{(q)}$. Denoting the eigenvalues and eigenvectors by $E_n^{(q)}$ and $\Psi_n^{(q)}$, i.e.,

$$H^{(q)}\Psi_n^{(q)} = E_n^{(q)}\Psi_n^{(q)}, \quad (12)$$

expressing $\Psi_n^{(q)}$ in the two-component form, and using Eq. (8), one can easily show that up to an undetermined scalar multiple, $\Psi_n^{(q)}$ has the following form:

$$\Psi_n^{(q)} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 - iqE_n^{(q)} \\ 1 + iqE_n^{(q)} \end{pmatrix} \Phi_n^{(q)}, \quad (13)$$

where $\Phi_n^{(q)} \in \mathcal{H}_t$ satisfies:

$$\left[\hat{D}_2 - iE_n^{(q)}(\hat{D}_1 - \frac{\dot{q}}{q}) - (E_n^{(q)})^2 \right] \Phi_n^{(q)} = 0. \quad (14)$$

This equation may be viewed as a ‘generalized’ eigenvalue equation¹ in $\mathcal{H}_t$. It defines both the vectors $\Phi_n^{(q)}$ and the complex numbers $E_n^{(q)}$. It reduces to the ordinary eigenvalue equation for $\hat{D}_2$, if $\hat{D}_1 = \dot{q}/q$. Note that this is also one of the conditions for self-adjointness of the Hamiltonian, with the choice of (11) for the inner product. Furthermore, if this condition is satisfied, then Eq. (14) determines $E_n^{(q)}$ up to a sign, i.e., eigenvalues come in pairs of opposite sign.

If q is chosen to be time-independent, then (14) does not carry any information about q and therefore $\Phi_n^{(q)}$ and $E_n^{(q)}$ are independent of the choice of q .² Hence, one can drop the labels (q) on the right hand side of Eq. (13). In this case, Eq. (14) becomes:

$$[\hat{D}_2 - iE_n\hat{D}_1 - E_n^2] \Phi_n = 0. \quad (15)$$

¹Note that this terminology has nothing to do with the concept of generalized eigenvalues of spectral analysis.

²This can also be seen by noting that under the transformation $\Psi_n^{(q)} \rightarrow \tilde{\Psi}_n^{(q)} = g(q', q)\Psi_n^{(q)}$, the eigenvectors preserve their form (13) and that $\tilde{\Psi}_n^{(q)}$ is an eigenvector of $\hat{H}^{(q')}$ with the same eigenvalue $E_n^{(q)}$.

Now let us use the inner product (11) to compute the inner product of two eigenvectors of the Hamiltonian. Performing the algebra, one finds

$$(\Psi_m^{(q)}, \Psi_n^{(q)}) = i(q^* E_m^* - q E_n) \langle \Phi_m | \Phi_n \rangle . \quad (16)$$

Therefore if q is a positive imaginary number, i.e., $q = i|q|$, then

$$(\Psi_m^{(q)}, \Psi_n^{(q)}) = |q|(E_m^* + E_n) \langle \Phi_m | \Phi_n \rangle . \quad (17)$$

Hence the eigenvectors $\Psi_n^{(q)}$ and $\Psi_m^{(q)}$ with $E_m = -E_n^*$ (if they exist) are orthogonal regardless of the value of $\langle \Phi_m | \Phi_n \rangle$. Furthermore, one has $(\Psi_n^{(q)}, \Psi_n^{(q)}) = 2|q|\text{Re}(E_n) \langle \Phi_n | \Phi_n \rangle$, i.e., the norm of an energy eigenvector has the same sign as the real part of the corresponding eigenvalue. It vanishes for the zero³ and imaginary energy eigenvalues. Note that here I am assuming that the inner product $\langle \cdot | \cdot \rangle$ on $\mathcal{H}_t$ is non-negative. In fact $\mathcal{H}_t$ is to be identified with the separable Hilbert space $L^2(\Sigma_t)$ of square-integrable functions on Σ_t where the integration is defined by the measure $[\det({}^{(3)}g)]^{1/2}$ defined by the Riemannian three-metric ${}^{(3)}g$ induced by the four-metric g .

Another interesting property of the inner product (11) is the fact that for imaginary q it yields the familiar Klein-Gordon inner product, $\langle \cdot, \cdot \rangle_{\text{KG}}$. This is easily seen by substituting (7) in (11), which leads to:

$$(\Psi_1, \Psi_2) = q \langle \Phi_1 | \dot{\Phi}_2 \rangle + q^* \langle \dot{\Phi}_1 | \Phi_2 \rangle = q [\langle \Phi_1 | \dot{\Phi}_2 \rangle - \langle \dot{\Phi}_1 | \Phi_2 \rangle] =: q \langle \Phi_1, \Phi_2 \rangle_{\text{KG}} . \quad (18)$$

It is also useful to recall that the space $\mathcal{H}_t \oplus \mathcal{H}_t$ is nothing but the space of the possible initial conditions $[\Phi(t, x^i), \dot{\Phi}(t, x^i)]$ with initial time being t and $(x^i) \in \Sigma_t$. In view of the well-posedness of the dynamical equation [25], this (vector) space is isomorphic to the space of solutions of the field equation (1). Hence a two-component decomposition may be viewed as a splitting of the space of solutions of the field equations. In view of the freedom of choice of the parameter q , this splitting is clearly not unique.

³The existence of a zero energy eigenvector $\Psi_0^{(q)}$ is equivalent to the existence of the solution to the differential equation $\hat{D}_2 \Phi_0 = 0$.

3 Cyclic States and Adiabatic Geometric Phase

By definition a cyclic state (an element of the projective Hilbert space) of a quantum mechanical system, whose dynamics is governed by the Schrödinger equation

$$i\dot{\psi}(t) = \hat{H}(t) \psi(t) , \quad (19)$$

is said to be cyclic with a period τ , if it is an eigenstate of the time-evolution operator $\hat{U}(\tau) := \mathcal{T} \exp[-i \int_0^\tau \hat{H}(t) dt]$. Here $\mathcal{T}$ is the time-ordering operator. An associated initial state vector $\psi(0)$ then satisfies:

$$\psi(\tau) = \hat{U}(\tau) \psi(0) = e^{i\alpha(\tau)} \psi(0) , \quad (20)$$

where $\alpha(\tau) \in \mathbb{C}$. If the Hamiltonian is self-adjoint, then $\alpha(\tau) \in \mathbb{R}$ and consequently $\psi(\tau)$ and $\psi(0)$ differ by a phase. In general $\alpha(\tau)$ may be expressed as the sum of a dynamical and a geometrical part [3]. This decomposition uses the inner product structure of the Hilbert space.

The situation is rather more transparent if the time-dependence of the Hamiltonian is adiabatic. In this case, one can follow Berry's approach [1] of employing the adiabatic theorem of quantum mechanics. According to the adiabatic theorem [26], if the initial state is an eigenstate of the initial Hamiltonian $\hat{H}(0)$, then after a time period $t > 0$ it evolves into an eigenstate of the Hamiltonian $\hat{H}(t)$. More precisely assume that the time-dependence of the Hamiltonian is realized through its dependence on a set of parameters $R = (R^1, \dots, R^n)$ and a smooth curve $C : [0, \tau] \rightarrow \mathcal{M}$, where R is viewed as coordinates of a parameter space $\mathcal{M}$, i.e., $\hat{H}(t) := H[R(t)]$, and $R(t) = (R^1(t), \dots, R^n(t)) := C(t)$. Furthermore, let $\psi_n[R]$ denote eigenvectors of $\hat{H}[R]$ with eigenvalue $E_n[R]$:

$$\hat{H}[R] \psi_n[R] = E_n[R] \psi_n[R] , \quad (21)$$

whose dependence on R is assumed to be smooth, the degree of the degeneracy of the eigenvalues is independent of R , and for $R = R(t)$ there is no level crossings. Then the statement of the adiabatic theorem may be summarized by the following (approximate) equation:

$$\psi(t) := \hat{U}(t) \psi_n(0) \approx e^{i\alpha_n(t)} \psi_n(t) , \quad (22)$$

where $\psi_n(t) := \psi_n[R(t)]$. If $E_n(t) := E_n[R(t)]$ is $\mathcal{N}$ -fold degenerate, then ψ_n belongs to the $\mathcal{N}$ -dimensional degeneracy subspace $\mathcal{H}_n$ and α_n is an $\mathcal{N} \times \mathcal{N}$ matrix-valued function of time. The approximation sign $\approx$ in (22) is used to remind one of the fact that this relation is only valid if the adiabatic approximation is justified.

Assuming the validity of the adiabatic approximation ($\approx \rightarrow =$) and substituting (22) in the Schrödinger equation (19), one has [2]:

$$e^{i\alpha_n(t)} = \exp[-i \int_0^t E_n(t') dt'] \mathcal{P} \exp[i \int_{C(0)}^{C(t)} \mathcal{A}_n] , \quad (23)$$

$$\mathcal{A}_n^{IJ}[R] := \frac{i \langle \psi_n^I[R], \frac{\partial}{\partial R^a} \psi_n^J[R] \rangle}{\langle \psi_n^I[R], \psi_n^I[R] \rangle} dR^a = \frac{i \langle \psi_n^I[R], d\psi_n^J[R] \rangle}{\langle \psi_n^I[R], \psi_n^I[R] \rangle} , \quad (24)$$

where $\mathcal{P}$ is the path-ordering operator, $\psi_n^I[R]$ form a complete orthonormal basis of the degeneracy subspace $\mathcal{H}_n$, and $\langle , \rangle$ is the inner product. If the Hamiltonian is periodic, i.e., C is a closed curve, then according to (22), $\psi_n[R(0)] = \psi_n[R(T)]$ is a cyclic state vector. In this case the first and the second (path-ordered) exponential in (23), with $t = T$, are called the dynamical and the geometrical parts of the total adiabatic matrix-valued phase $\exp[i\alpha_n(T)]$, respectively, [2]. The qualification ‘geometrical’ is best justified by identifying the geometric part of the phase by the holonomy of a principal spectral bundle over the parameter space $\mathcal{M}$ or alternatively the universal classifying bundle over the projective Hilbert space, [27, 3, 28].

The situation for a non-self-adjoint Hamiltonian is rather more complicated. The dynamical and the geometrical phase can still be defined in terms of the projective Hilbert space [29]. However, in general the eigenvectors of the Hamiltonian are not orthogonal.⁴ This renders the proof of the adiabatic theorem [26] invalid. One can still postulate (22) as an ansatz which may or may not be valid for specific evolutions. The condition of the validity of this ansatz, which allows one to pursue the same strategy in defining the adiabatic geometric phase, is

$$\langle \psi_m, \psi_n \rangle \langle \dot{\psi}_n, \dot{\psi}_n \rangle = \langle \dot{\psi}_m, \dot{\psi}_n \rangle \langle \psi_n, \psi_n \rangle , \quad (25)$$

where ψ_m and ψ_n are any pair of distinct eigenvectors of the Hamiltonian. This condition is obtained by substituting the ansatz (22) in the Schrödinger equation (19) and taking the inner

⁴Note that the eigenvectors within a single degeneracy subspace can always be orthonormalized. However the eigenvectors corresponding to distinct eigenvalues in general overlap.

product of both sides of the resulting equation with ψ_m . Eq. (25) is trivially satisfied for the case of a self-adjoint Hamiltonian. In this case, the left hand side vanishes identically and $\langle \dot{\psi}_m, \dot{\psi}_n \rangle$ for $m \neq n$, vanishes approximately by virtue of the adiabatic approximation [30].

Before pursuing the derivation of the expression for the geometric phase, I must emphasize that a general cyclic two-component state vector is clearly cyclic in its both components. Identifying the corresponding function space $\mathcal{H}_t \oplus \mathcal{H}_t$ (Note that the t -dependence is only relevant to the inner product structure and the vector space structure is independent of t .) with the space of all possible initial data, a cyclic two-component state vector $\Psi^{(q)}(0)$ which by definition satisfies $\Psi^{(q)}(\tau) = \exp[i\alpha(\tau)]\Psi^{(q)}(0)$, is associated with a ‘cyclic’ solution of the Klein-Gordon equation (1) whose velocity is also cyclic with the same (possibly non-unimodular) phase and period, i.e., $\Phi(\tau, x^i) = \exp[i\alpha(\tau)]\Phi(0, x^i)$ and $\dot{\Phi}(\tau, x^i) = \exp[i\alpha(\tau)]\dot{\Phi}(0, x^i)$. This is in contrast with the usual definition of cyclicity for classical fields [4, 5]. It seems more reasonable to ascribe the term ‘cyclic’ to a repetition, up to a scalar multiple, of all initial conditions rather than to just one of them. In this article I shall use the term *cyclic* in this sense.

Next, let us consider the two-component formulation of the Klein-Gordon equation. Suppose for simplicity that $E_n^{(q)}$ of Eq. (12) is independent of q , i.e., $E_n^{(q)} = E_n$ and that it is non-degenerate. Then, a direct generalization of the concept of adiabatic evolution in non-relativistic quantum mechanics suggests one to use the ansatz:

$$\Psi^{(q)}(t) \approx e^{i\alpha_n(t)} \Psi_n^{(q)}[R(t)] , \quad (26)$$

to define the relativistic analog of adiabatic evolution. One can show, however, that this ansatz leads to a restrictive notion of adiabatic approximation. I shall refer to this approximation as the *ultra-adiabatic approximation*. More precisely, I shall use the following definition:

Definition 1: A two-component state vector $\Psi^{(q)}(t)$ is said to undergo an *exact ultra-adiabatic evolution* if and only if

$$\Psi^{(q)}(t) = e^{i\alpha_n(t)} \Psi_n^{(q)}[R(t)] , \quad (27)$$

for some n and α_n .

Note that Definition 1 also provides a definition for *ultra-adiabatic approximation* by replacing Eq. (27) by Eq. (26).

In order to derive the conditions under which the ultra-adiabatic approximation is valid, one must substitute Eqs. (27), (12), and (13) in the Schrödinger equation (5). This yields

$$\left[-\dot{\alpha}_n(1 - iqE_n) + q\dot{E}_n - E_n(1 - iqE_n) \right] \Phi_n + i(1 - iqE_n)\dot{\Phi}_n = 0 , \quad (28)$$

$$\left[-\dot{\alpha}_n(1 + iqE_n) - q\dot{E}_n - E_n(1 + iqE_n) \right] \Phi_n + i(1 + iqE_n)\dot{\Phi}_n = 0 . \quad (29)$$

Adding both sides of these equations and simplifying the result, one has

$$(\dot{\alpha}_n + E_n)\Phi_n - i\dot{\Phi}_n = 0 . \quad (30)$$

This equation leads directly to the expression for the total phase (23) with the Berry connection one-form given by

$$\mathcal{A}_n = \frac{i\langle \Phi_n | \frac{\partial}{\partial R^a} \Phi_n \rangle}{\langle \Phi_n | \Phi_n \rangle} dR^a = \frac{i\langle \Phi_n | d\Phi_n \rangle}{\langle \Phi_n | \Phi_n \rangle} . \quad (31)$$

Here R denotes the parameters of the system, i.e., the metric g , the electromagnetic potential A and the scalar potential V . Moreover I have used the identity $\dot{\Phi}_n dt = (\partial \Phi_n / \partial R^a) dR^a = d\Phi_n$.

Furthermore, subtracting Eq. (29) from (28) and using Eq. (30) to simplify the resulting expression, one finds

$$\dot{E}_n = 0 . \quad (32)$$

This condition which is a direct consequence of Definition 1 does not have a counterpart in ordinary non-relativistic quantum mechanics. Its roots may be sought in the fundamental difference between ordinary (one-component) Schrödinger and Klein-Gordon equations. One may argue based on physical grounds that the condition (32) and consequently the concept of the ultra-adiabatic evolution are too restrictive. Indeed it is possible to relax this condition by adopting a more general definition of adiabatic evolution. For the moment, however, I shall continue with a further analysis of the ultra-adiabatic evolutions.

Because Eq. (30) is identical with the one obtained in the non-relativistic case, in addition to condition (32) one also has the analog of Eq. (25). If Φ_n turn out to be orthogonal, the latter reduces to

$$\langle \Phi_m | \dot{\Phi}_n \rangle = 0 , \quad \forall m \neq n , \quad (33)$$

which is the well-known condition for the exactness of the adiabatic approximation in non-relativistic quantum mechanics, [30]. Hence, for the cases where Φ_n are orthogonal, the ultra-adiabatic approximation is exact if and only if Eqs. (32) and (33) are satisfied.

Note that Eqs. (28) and (29) and consequently conditions (32) and (33) are valid if and only if the ultra-adiabatic approximation is exact. As one knows from non-relativistic quantum mechanics, the condition of the exactness of (ultra-)adiabatic approximation is highly restrictive.⁵ More interesting are cases where the (ultra-)adiabatic approximation is valid only approximately, i.e., cases where instead of (27), (26) holds. In this case, Eqs. (28), (29), and conditions (32) and (33) are required to be satisfied approximately, namely

$$\dot{E}_n \approx 0, \quad (34)$$

$$\langle \Phi_m | \dot{\Phi}_n \rangle \approx 0, \quad \forall m \neq n. \quad (35)$$

More precisely, the ultra-adiabatic approximation is a valid approximation if and only if (34) and (35) are satisfied. The precise meaning of the $\approx$ in these equations will be clarified momentarily.

In the above discussion, the condition of time-independence of q does not play any significant role in the derivation of Eqs. (30) and (31). In fact, allowing q to be time-dependent only changes the term $q\dot{E}_n$ in Eqs. (28) and (29) to $d(qE_n^{(q)})/dt$. Therefore, up on adding the resulting equations one still obtains Eq. (30). The only consequences of using a time-dependent q are the emergence of q -dependent E_n and Φ_n and the condition

$$\frac{d}{dt}(qE_n^{(q)}) \approx 0, \quad (36)$$

which generalizes (34).

There is a particular case in which q may be time-dependent but E_n and Φ_n are still independent of the choice of q . This is the case, where the operator $\hat{D}_1$ of (3) is zero-th order and it only involves time-dependent functions. In this case one can choose q in such a way as to satisfy $\hat{D}_1 = \dot{q}/q$. This condition reduces Eq. (14) to the eigenvalue equation for $\hat{D}_2$, with eigenvalues E_n^2 and eigenvectors Φ_n . Hence, E_n and Φ_n are still q -independent. I shall show in section 6 how this apparently very special case may be realized and used in the study of spatially homogeneous (Bianchi) cosmological models.

⁵In fact, the (ultra-)adiabatic approximation is exact if and only if the evolving state is stationary [31].

The appearance of the decomposition parameter q in (36) and the fact that this condition has no non-relativistic analog suggests that perhaps the notion of ultra-adiabatic approximation is too limited. In order to obtain a physically more appealing concept of adiabatic approximation, one must consider a more general ansatz than (26).

Consider the general solutions Ψ of the two-component Schrödinger equation (5) of the form

$$\Psi^{(q)} = \sum_n e^{i\alpha_n} \Psi_n^{(q)} , \quad (37)$$

where $\alpha_n \in \mathbb{C}$ and $\Psi_n^{(q)}$ are the eigenvectors of the two-component Hamiltonian (8). Substituting Eq. (37) in the Schrödinger equation (5) and making use of Eqs. (12) and (13), one has

$$\sum_n e^{i\alpha_n} \left\{ [(E_n^{(q)} + \dot{\alpha}_n)(1 + iqE_n^{(q)}) + \frac{d}{dt}(qE_n^{(q)})] \Phi_n^{(q)} - i(1 - iqE_n^{(q)}) \dot{\Phi}_n^{(q)} \right\} = 0 , \quad (38)$$

$$\sum_n e^{i\alpha_n} \left\{ [(E_n^{(q)} + \dot{\alpha}_n)(1 - iqE_n^{(q)}) - \frac{d}{dt}(qE_n^{(q)})] \Phi_n^{(q)} - i(1 + iqE_n^{(q)}) \dot{\Phi}_n^{(q)} \right\} = 0 . \quad (39)$$

Adding and subtracting both sides of these equations and simplifying the result lead to

$$\sum_n e^{i\alpha_n} [(E_n^{(q)} + \dot{\alpha}_n) \Phi_n^{(q)} - i \dot{\Phi}_n^{(q)}] = 0 , \quad (40)$$

$$\sum_n e^{i\alpha_n} \left\{ \left[\frac{d}{dt}(qE_n^{(q)}) \right] \Phi_n^{(q)} + iqE_n^{(q)} [(E_n^{(q)} + \dot{\alpha}_n) \Phi_n^{(q)} - i \dot{\Phi}_n^{(q)}] \right\} = 0 . \quad (41)$$

Next assume that $\hat{D}_2$ is a non-degenerate self-adjoint operator with a discrete spectrum and $\hat{D}_1 = \dot{q}/q$. Then, E_n and Φ_n do not depend on q and $\hat{D}_2 \Phi_n = E_n^2 \Phi_n$. Now, differentiate both sides of the latter equation with respect to time and take their inner product with Φ_m . Since in this case Φ_n are orthogonal, one has the well-known identity [1]

$$\frac{\langle \Phi_m | \dot{\hat{D}}_2 | \Phi_n \rangle}{E_n^2 - E_m^2} = \langle \Phi_m | \dot{\Phi}_n \rangle , \quad \text{for all } m \neq n , \quad (42)$$

where Φ_n and Φ_m correspond to distinct eigenvalues of $\hat{D}_2$, i.e., $E_n^2 \neq E_m^2$.

Quantum adiabatic approximation is valid if the left hand side of this equation which involves the time-derivative of $\hat{D}_2$ can be neglected, [30]. This statement provides the true meaning of the condition (35)

$$\langle \Phi_m | \dot{\Phi}_n \rangle = \frac{\langle \Phi_m | \dot{\hat{D}}_2 | \Phi_n \rangle}{E_n^2 - E_m^2} \approx 0 , \quad \text{for all } m \neq n , \quad (43)$$

for the case where the above assumptions are valid. I shall next use this condition to define the notion of *adiabatic evolution* in relativistic scalar quantum mechanics.

For convenience, I shall use the notation Ψ_{-n} for the two-component eigenvector corresponding to the eigenvalue $E_{-n} := -E_n$. Since for each pair $(-n, n)$ there is a single Φ_n , one can write Eqs. (40) and (41) in the form

$$\sum_{n \geq 0} \left\{ \left[E_n(e^{i\alpha_n} - e^{i\alpha_{-n}}) + (\dot{\alpha}_n e^{i\alpha_n} + \dot{\alpha}_{-n} e^{i\alpha_{-n}}) \right] \Phi_n - i(e^{i\alpha_n} + e^{i\alpha_{-n}}) \dot{\Phi}_n \right\} = 0, \quad (44)$$

$$\sum_{n \geq 0} \left\{ \left[(e^{i\alpha_n} - e^{i\alpha_{-n}}) \frac{d}{dt}(qE_n) + iqE_n^2(e^{i\alpha_n} + e^{i\alpha_{-n}}) + iqE_n(\dot{\alpha}_n e^{i\alpha_n} - \dot{\alpha}_{-n} e^{i\alpha_{-n}}) \right] + qE_n(e^{i\alpha_n} - e^{i\alpha_{-n}}) \dot{\Phi}_n \right\} = 0. \quad (45)$$

Enforcing condition (43), one can reduce (44) and (45) to

$$E_n(e^{i\alpha_n} - e^{i\alpha_{-n}}) + (\dot{\alpha}_n e^{i\alpha_n} + \dot{\alpha}_{-n} e^{i\alpha_{-n}}) - (e^{i\alpha_n} + e^{i\alpha_{-n}}) a_n \approx 0, \quad (46)$$

$$(-if_n - a_n)(e^{i\alpha_n} - e^{i\alpha_{-n}}) + E_n(e^{i\alpha_n} + e^{i\alpha_{-n}}) + \dot{\alpha}_n e^{i\alpha_n} - \dot{\alpha}_{-n} e^{i\alpha_{-n}} \approx 0, \quad (47)$$

where $n \geq 0$ and

$$a_n := \frac{i\langle \Phi_n | \dot{\Phi}_n \rangle}{\langle \Phi_n | \Phi_n \rangle}, \quad f_n := \frac{\frac{d}{dt}(qE_n)}{qE_n} = \frac{d}{dt} \ln(qE_n).$$

Adding and subtracting both sides of (46) and (47) and assuming that $e^{i\alpha_n}$ is not negligibly small, one finds

$$-if_n(1 - e^{-i(\alpha_n - \alpha_{-n})}) + 2(E_n + \dot{\alpha}_n - a_n) \approx 0, \quad (48)$$

$$-if_n(e^{i(\alpha_n - \alpha_{-n})} - 1) + 2(E_n - \dot{\alpha}_{-n} + a_n) \approx 0. \quad (49)$$

Next, define $\eta_n^- := \alpha_n - \alpha_{-n}$, add both sides of (48) and (49), and simplify the result. This leads to

$$\dot{\eta}_n^- + f_n \sin \eta_n^- + 2E_n \approx 0. \quad (50)$$

Introducing $\eta_n^+ := \alpha_n + \alpha_{-n}$ and using (50), one can then express (48) in the form

$$\dot{\eta}_n^+ - if_n(1 - \cos \eta_n^-) - 2a_n \approx 0, \quad (51)$$

Hence in view of the definition $\eta_n^\pm := \alpha_n \pm \alpha_{-n}$, one has

$$\alpha_{\pm n} = \frac{1}{2}(\eta_n^+ \pm \eta_n^-) \approx \gamma_n + \delta_{\pm n}, \quad \forall n \geq 0, \quad (52)$$

$$\gamma_n := \int_0^t a_n(t') dt' = \int_{R(0)}^{R(t)} \mathcal{A}_n[R] , \quad (53)$$

$$\delta_{\pm n} := \frac{i}{2} \int_0^t f_n(t') [1 - \cos \eta_n(t')] dt' \pm \frac{\eta_n(t)}{2} , \quad (54)$$

where I have used Eq. (31) and η_n is the solution of

$$\dot{\eta}_n + f_n \sin \eta_n + 2E_n = 0 , \quad \text{with} \quad \eta_n(0) = 0 . \quad (55)$$

As seen from (52)-(54), the part γ_n of $\alpha_{\pm n}$ which is independent of E_n has the same form as the geometric phase angle of the non-relativistic quantum mechanics. In contrast, the part $\delta_{\pm n}$ of $\alpha_{\pm n}$ which does depend on $E_{\pm n}$ and plays the role of the dynamical phase angle, has a different expression from its non-relativistic counterpart. For the case of an ultra-adiabatic evolution where condition (36) is satisfied, $f_n \approx 0$ and

$$\delta_{\pm n} = \mp \int_0^t E_n(t') dt' = - \int_0^t E_{\pm n}(t') dt' .$$

This is identical with the expression for the non-relativistic adiabatic dynamical phase.

The above analysis shows that taking (43) as the defining condition for the *adiabatic approximation*, one obtains the same expression for the geometric phase as in the ultra-adiabatic case. This condition modifies the expression for the dynamical phase. In fact, the dynamical phase angle splits into a pair (δ_{-n}, δ_n) of dynamical angles. The latter is a consequence of the violation of the ultra-adiabaticity condition (43).

The relativistic adiabatic approximation outlined in the preceding paragraphs corresponds to the following definition of relativistic adiabatic evolution

Definition 2: A two-component state vector $\Psi^{(q)}(t)$ is said to undergo an *exact adiabatic evolution* if and only if

$$\Psi^{(q)}(t) = e^{i\alpha_n(t)} \Psi_n^{(q)}[R(t)] + e^{i\alpha_{-n}(t)} \Psi_{-n}^{(q)}[R(t)] , \quad (56)$$

for some n and $\alpha_{\pm n}$.

The *approximate adiabatic approximation* corresponds to the case where (56) is approximately valid, i.e.,

$$\Psi^{(q)}(t) \approx e^{i\alpha_n(t)} \Psi_n^{(q)}[R(t)] + e^{i\alpha_{-n}(t)} \Psi_{-n}^{(q)}[R(t)] . \quad (57)$$

This approximation is valid if and only if $\langle \Phi_m | \dot{\Phi}_n \rangle \approx 0$ for all $m \neq n$. Definition 2 provides a suitable definition for an adiabatic evolution in relativistic (scalar) quantum mechanics. In particular, it ensures that for a cyclic change of the parameters of the system, the one-component Klein-Gordon field and its time-derivative have cyclic evolutions. The difference between the ultra-adiabatic and adiabatic evolutions is that for a cyclic ultra-adiabatic evolution the (possibly non-unimodular) phases of the one-component field and its time-derivative are required to be equal, whereas in a cyclic adiabatic evolution these phases are generally different.

Since the defining condition for the relativistic and non-relativistic adiabatic evolution are identical, one can use the well-known results of non-relativistic quantum mechanics to generalize the above results to the case where E_n is degenerate.

An important aspect of the above derivation of the geometric phase is that it does not use the particular form of an inner product on $\mathcal{H}_t \oplus \mathcal{H}_t$, i.e., the Hermitian matrix h of (10). It only uses the inner product on $\mathcal{H}_t$. One might argue that based on the known features of the non-relativistic case, the independence of the geometric phase from the inner product is quite natural and thus not particularly important. A review of the existing literature [6, 17] shows, however, that in the previously studied examples a great deal of effort was made to define an inner product on the space of solutions before the problem of the geometric phase could be addressed. The construction of such an inner product is a highly technical problem and a satisfactory solution for arbitrary spacetimes is not known. The results of this section indicates that indeed one does not need to construct an inner product on the space of solutions. What is needed is the L^2 inner product on $\mathcal{H}_t$ which is naturally given by the induced spatial metric. In this way, one can conveniently avoid the difficult problem of constructing an inner product on the space of solutions and carry on with the analysis of the adiabatic geometric phase.

In the following sections the practical advantages of the two-component formulation are demonstrated for three physical examples.

4 Rotating Magnetic Field in Minkowski Background

Consider the geometric phase induced on a Klein-Gordon field in a Minkowski background due to a rotating magnetic field. This problem was originally studied by Anandan and Mazur [6] using the one-component formalism.

In this case, in a global cartesian coordinate system, one has $g_{00} = -1$, $g_{ij} = \delta_{ij}$, $g_{0i} = V = 0$, and $\mathcal{H}_t = L^2(\mathbb{R}^3)$. Following [6], let us first consider the case of a constant magnetic field along the x^3 -axis. Then in the symmetric gauge, one has $A_0 = A_3 = 0$, $A_1 = -Bx^2/2$, and $A_2 = Bx^1/2$. Substituting these equations in Eqs. (3) and (4), one finds $\hat{D}_1 = 0$ and

$$\hat{D}_2 = -\nabla^2 - ieB \frac{\partial}{\partial \varphi} + \frac{e^2 B^2}{4} \rho^2 + \mu^2, \quad (58)$$

where ∇^2 is the Laplacian and (ρ, φ, x^3) are cylindrical coordinates in $\mathbb{R}^3$. Clearly $\hat{D}_2$ is self-adjoint. Therefore, Eq. (14) reduces to the eigenvalue equation for $\hat{D}_2$, namely Φ_n are orthogonal eigenvectors of $\hat{D}_2$ with eigenvalue E_n^2 . Taking $q = i$ in Eq. (7) and choosing the inner product (11), the Hamiltonian $H^{(i)}$ of (5) is also self-adjoint.

The situation is quite similar to the non-relativistic Landau level problem. Clearly, Φ_n are infinitely degenerate. They are given by

$$\Phi_n^{(p,m)} = N_n e^{ipx^3} e^{im\varphi} \chi_{nmp}(\rho), \quad (59)$$

where $p \in \mathbb{R}$, $m = 0, 1, 2, \dots$ label the vectors within the degeneracy subspace $\mathcal{H}_n$, χ_{nmp} are orthogonal solutions of

$$\left[\frac{d^2}{d\rho^2} + \frac{1}{\rho} \frac{d}{d\rho} + (k^2 - \frac{m^2}{\rho^2} - \lambda^2 \rho^2) \right] \chi_{nmp}(\rho) = 0, \quad (60)$$

$$k^2 := E_n^2 - (p^2 + \mu^2 + emB), \quad \lambda := \frac{eB}{2}$$

and N_n are normalization constants chosen in such a way as to ensure

$$\langle \Phi_{\tilde{n}}^{(\tilde{p}, \tilde{m})} | \Phi_n^{(pm)} \rangle = \delta(\tilde{n}, n) \delta(\tilde{m}, m) \delta(\tilde{p}, p). \quad (61)$$

Here $\delta(,)$ denotes a Kronecker or a Dirac delta function depending on whether the arguments are discrete or continuous, respectively.

In order to solve the eigenvalue problem for the rotating magnetic field, one can easily use the unitary transformations [6, 31]

$$\mathcal{U}(\theta, \varphi) = e^{-i\varphi\hat{J}_3} e^{-i\theta\hat{J}_2} e^{i\varphi\hat{J}_3} , \quad (62)$$

relating the eigenvectors Φ_n of $\hat{D}_2$ to those corresponding to the constant magnetic field (59). In Eq. (62), θ and φ are azimuthal and polar angles in spherical coordinates and $\hat{J}_i$ are angular momentum operators (generators of $SO(3)$) acting on the Hilbert space $L^2(\mathbb{R}^3)$. $\mathcal{U}(\theta, \varphi)$ are well-defined everywhere except along the negative x^3 -axis which can be excluded by assuming that $\vec{B}(t) = (B, \theta(t), \varphi(t))$ does not cross this axis. Otherwise, one may choose another coordinate frame and remedy the problem by performing appropriate gauge transformations as described in Ref. [31] for the non-relativistic case. Clearly,

$$\begin{aligned} \hat{D}_2[\vec{B}(t)] &= \mathcal{U}(\theta(t), \varphi(t)) \hat{D}_2[\vec{B} = B\hat{x}^3] \mathcal{U}^\dagger(\theta(t), \varphi(t)) , \\ \Phi_n[\vec{B}(t)] &= \mathcal{U}(\theta(t), \varphi(t)) \Phi_n[\vec{B} = B\hat{x}^3] , \\ E_n[\vec{B}(t)] &= E_n[\vec{B} = B\hat{x}^3] = \text{constant} . \end{aligned}$$

The latter relation which implies $\dot{E}_n = 0$ indicates that an adiabatic evolution of this system is, in fact, ultra-adiabatic.

As noted in Ref. [6], the presence of the degeneracy leads to non-Abelian geometric phases (23) defined by the connection one-form $\mathcal{A}_n$, (24). The components of $\mathcal{A}_n$ are given by the non-Abelian generalization of (31), namely

$$A_n^{IJ} = i\langle \Phi_n^{(I)} | d\Phi_n^{(J)} \rangle , \quad (63)$$

and are independent of the choice of the matrix h of (10). In Eq. (63), $I := (p, m)$ and $J := (p', m')$, and use is made of (61). I shall not be elaborating on this problem any further since the specific results are exactly the same as the ones reported in Ref. [6]. It is however worth mentioning that each Φ_n defines a pair of orthonormal two-component eigenvectors $\Psi_{\pm n}^{(i)}$ corresponding to the choices $\pm E_n$ for each eigenvalue E_n^2 of $\hat{D}_2$. Hence in this case the two-component formalism reproduces the results of [6] which were obtained using a more subtle method of taking square root of the second order Klein-Gordon operator and projecting onto the spaces of negative and positive energy (frequency) solutions of the Klein-Gordon equation.

5 Rotating Cosmic String

In Ref. [17], the authors study the geometric (or rather topological) phases induced on a Klein-Gordon field due to a rotating cosmic string. In this section, I shall outline a solution to this problem using the two-component formalism.

The local coordinate expression for the metric corresponding to a rotating cosmic string with angular momentum j and linear mass density d is [17]:

$$g = \begin{pmatrix} -1 & 0 & -4j & 0 \\ 0 & 1 & 0 & 0 \\ -4j & 0 & (\alpha\rho)^2 - (4j)^2 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad (64)$$

where $(x^\mu) = (t, \rho, \varphi, z)$ and (ρ, φ, z) are cylindrical coordinates on the spatial hypersurface Σ_t and $\alpha := 1 - 4d$. Σ_t corresponds to a cone with a deficit angle $\beta = 8\pi d = 2\pi(1 - \alpha)$.

Note that for $\rho \leq 4j/\alpha$, $\partial/\partial\varphi$ becomes timelike. This leads to the existence of closed timelike curves. This region can be ignored by imposing appropriate boundary conditions on the fields, i.e., $\Phi = 0$ for $\rho \leq 4j/\alpha$.

Performing the necessary calculations, one finds the following expressions for the operators $\hat{D}_1$ and $\hat{D}_2$ of Eqs. (3) and (4):

$$\hat{D}_1 = \frac{8j}{(\alpha\rho)^2 - (4j)^2} \frac{\partial}{\partial\varphi}, \quad (65)$$

$$\hat{D}_2 = \left(\frac{-1}{1 - (\frac{4j}{\alpha\rho})^2} \right) \left[\frac{\partial^2}{\partial\rho^2} + \frac{1}{\rho} \frac{\partial}{\partial\rho} + \frac{1}{(\alpha\rho)^2} \frac{\partial^2}{\partial\varphi^2} + \frac{\partial^2}{\partial z^2} - \mu^2 \right]. \quad (66)$$

Therefore the conditions for the self-adjointness of the Hamiltonian $H^{(a)}$ of (5) cannot be met. Let us proceed, however, with considering the eigenvectors of $\Psi_n^{(a)}$ of $H^{(a)}$, (12). For the metric (64), Eq. (15) takes the form:

$$\left\{ \frac{\partial^2}{\partial\rho^2} + \frac{1}{\rho} \frac{\partial}{\partial\rho} + \frac{1}{(\alpha\rho)^2} \frac{\partial^2}{\partial\varphi^2} + \frac{i8jE_n}{(\alpha\rho)^2} \frac{\partial}{\partial\varphi} + \frac{\partial^2}{\partial z^2} - \mu^2 + [1 - (\frac{4j}{\alpha\rho})^2]E_n^2 \right\} \Phi_n = 0. \quad (67)$$

In view of an observation made in Ref. [18] and used in [17], let us write Φ_n in the form $\Phi_n = \exp(i\zeta\varphi)\phi_n$. Substituting this equation in (67), one finds that for $\zeta = -4jE_n$, ϕ_n satisfies:

$$\left\{ \frac{\partial^2}{\partial\rho^2} + \frac{1}{\rho} \frac{\partial}{\partial\rho} + \frac{1}{(\alpha\rho)^2} \frac{\partial^2}{\partial\varphi^2} + \frac{\partial^2}{\partial z^2} - \mu^2 + E_n^2 \right\} \phi_n = 0. \quad (68)$$

Eq. (68) may be obtained from (67) by setting $j = 0$ and replacing Φ_n by ϕ_n . Hence ϕ_n determine the eigenvectors of the Hamiltonian for a non-rotating string of the same mass density. In this case $\hat{D}_1$ vanishes and $\hat{D}_2$ becomes self-adjoint. Therefore, ϕ_n are orthogonal eigenvectors of $\hat{D}_2$, with $j = 0$. Choosing $q = i$ and adopting the inner product (11), the corresponding Hamiltonian becomes also self-adjoint.

In fact, it is not difficult to show that the solutions of Eq. (68) are of the form:

$$\phi_n = N_n e^{ipz} e^{im\varphi} J_\nu(k\rho) , \quad (69)$$

where N_n are appropriate normalization constants, J_ν are Bessel functions, and

$$k := \sqrt{E_n^2 - (p^2 + \mu^2)} , \quad \nu := m/\alpha .$$

The orthogonality property of ϕ_n carries over to Φ_n since the measure of the integration on Σ_t is independent of j . This is because of the identity:

$$\det[g] = -\det[{}^{(3)}g] ,$$

which holds for any metric with the lapse function $N = 1$, [25].

The situation is analogous to the case of a rotating magnetic field. However, in this case the Klein-Gordon field acquires an Aharonov-Bohm type phase which is topological in nature. As Berry describes in his (by now classic) article [1], the Aharonov-Bohm phase may be viewed as a particular case of a geometric phase. This is done, for the original Aharonov-Bohm system of an electron encircling a confined magnetic flux line, by considering the electron to be localized in a box which is then carried around the flux line. Thus the time-dependence of the system is introduced by choosing a coordinate system centered inside the box. This leads to geometric phases for the energy eigenfunctions. The same result is then applied to the electron wave packet, only because the geometric phase is independent of the energy eigenvalues, i.e., all the energy eigenvectors and therefore any linear combinations of them, in particular the one forming the localized electron wave packet, acquire the same geometric phase which is then shown to be the same as the one discovered by Aharonov and Bohm [32].

Ref. [17] uses the analogy between the system of rotating cosmic string and that of Aharonov and Bohm to obtain the corresponding geometric phases. This is however not quite justified

for arbitrary energy eigenfunctions since as shown below and also in [17], unlike the Aharonov-Bohm system, the induced phase in this case does depend on the energy eigenvalue. Consequently an arbitrary localized Klein-Gordon field which is a superposition of different energy eigenfunctions will not be cyclic. Berry's argument therefore applies only to those localized field configurations which are energy eigenfunctions. Such eigenfunctions do actually exist. Particular examples can be constructed by virtue of the infinite degeneracy arising from the axial symmetry of the problem. This is demonstrated in Appendix A.

Therefore, we can proceed with using the analogy with Berry's treatment of the Aharonov-Bohm phase [1] to derive the geometric phase in the framework of the two-component formalism. This is done by changing to a frame centered in a box which circulates around the string at a distance larger than $4j/\alpha$. If R^i are coordinates of the center of the box and x'^i are the coordinates centered at $R = (R^i)$, then the eigenfunctions are of the form: $\Phi_n(x') = \Phi_n(x - R)$. Substituting this expression in the non-Abelian version of (31), one finds

$$\begin{aligned} \mathcal{A}_n^{IJ} &= i \langle \Phi_n^I(x - R) | \frac{\partial}{\partial R^i} | \Phi_n^J(x - R) \rangle dR^i, \\ &= i \int_{\Sigma} d\Omega \phi_n^{*I}(x - R) \left[-4i E_n j \phi_n^J(x - R) dR^2 + \frac{\partial}{\partial R^i} \phi_n^J(x - R) dR^i \right], \\ &= 4j E_n \delta_{IJ} dR^2, \end{aligned} \tag{70}$$

where $d\Omega = \alpha \rho d\rho d\varphi dz$, I and J stand for possible degeneracy labels corresponding to localized eigenfunctions, R^2 is the polar angle associated with the center of the box, and ϕ_n are assumed to be normalized. For a curve C with winding number N_C , the geometric phase 'angle' is given by

$$\gamma_n = N_C \int_0^{2\pi} \mathcal{A}_n = 8\pi j E_n N_C, \tag{71}$$

where the labels I, J and δ_{IJ} have been suppressed for convenience. This is identical with the result of Ref. [17]. Note however that here I have not been concerned with the consideration of the difficult problem of the choice of an inner product for the space of the solutions of the Klein-Gordon equation $(\mathcal{H}_t \oplus \mathcal{H}_t)$, such as the one proposed by Ashtekar and Mangon [33] and apparently 'used' by Corichi and Pierri in Ref. [17]. In fact, as I have shown in section 3, the geometric phase is independent of the particular choice of such an inner product. This is

also implicit in the Corichi and Pierri's derivation of the geometric phase in [17]. Although they discuss the Ashtekar-Mangon scheme in some detail, the final derivation does not use the particular form of the inner product.

It is also worth mentioning that although the eigenvalues E_n may be degenerate, the corresponding geometric phase is still Abelian.

6 Spatially Homogeneous Cosmological Models

Consider Klein-Gordon fields in a spatially homogeneous (Bianchi) cosmological background associated with a Lie group G , i.e., $M = \mathbb{R} \times G$. In a synchronous invariant basis the spacetime metric g is given by its spatial components g_{ab} :

$$ds^2 = g_{\alpha\beta}\omega^\alpha\omega^\beta = -dt^2 + g_{ab}\omega^a\omega^b, \quad (72)$$

where ω^a are the left invariant one-forms and $g_{ab} = g_{ab}(t)$. Throughout this article I use the conventions of Ref. [34].

The first step in the study of the phenomenon of geometric phase due to a spatially homogeneous cosmological background is to compute the operators $\hat{D}_1$ and $\hat{D}_2$ of Eqs. (3) and (4) in the invariant basis. It is not difficult to see that with some care these equations are valid for arbitrary basis. One must only replace the coordinate labels $(\mu, \nu, \dots, i, j, \dots)$ with the basis (in this case invariant basis) labels $(\alpha, \beta, \dots, a, b, \dots)$, and interpret ∂_a as the action of the operators $\hat{X}_a$ associated with the dual vector fields to ω^a . This leads to:

$$\hat{D}_1 = g^{ab}\Gamma_{ab}^0, \quad (73)$$

$$\hat{D}_2 = -\Delta_t + \mu^2, \quad (74)$$

$$\Delta_t := g^{ab}\nabla_a\nabla_b = g^{ab}\hat{X}_a\hat{X}_b - \Gamma_{ab}^c\hat{X}_c, \quad (75)$$

where Δ_t is the Laplacian on Σ_t , ∇_a are the covariant derivatives corresponding to the Levi Civita connection, and [34]:

$$\Gamma_{\alpha\beta}^\gamma := \frac{1}{2}g^{\gamma\delta}(g_{\delta\alpha,\beta} + g_{\beta\delta,\alpha} - g_{\alpha\beta,\delta} + g_{\epsilon\alpha}C_{\delta\beta}^\epsilon + g_{\epsilon\beta}C_{\delta\alpha}^\epsilon) - \frac{1}{2}C_{\alpha\beta}^\gamma. \quad (76)$$

Here $g_{\alpha\beta,\gamma} = \hat{X}_\gamma g_{\alpha\beta}$ and $C_{\alpha\beta}^\gamma$ are structure constants:

$$[\hat{X}_\alpha, \hat{X}_\beta] = -C_{\alpha\beta}^\gamma \hat{X}_\gamma, \quad (77)$$

with $\hat{X}_0 := \partial/\partial t$. In view of the latter equality, the structure constants with a time label vanish. This simplifies the calculations of Γ 's. The only nonvanishing ones are

$$\Gamma_{ab}^0 = \frac{1}{2} \dot{g}_{ab}, \quad (78)$$

$$\Gamma_{ab}^c = \frac{1}{2} g^{cd} (g_{ea} C_{db}^e + g_{eb} C_{da}^e) - \frac{1}{2} C_{ab}^c. \quad (79)$$

In view of these relations, the expression for $\hat{D}_1$ and $\hat{D}_2$ may be further simplified:

$$\hat{D}_1 = \frac{\partial}{\partial t} \ln \sqrt{g}, \quad (80)$$

$$\hat{D}_2 = -\Delta_t + \mu^2 = -(g^{ab} \hat{X}_a \hat{X}_b - C_{ab}^b g^{ac} \hat{X}_c) + \mu^2, \quad (81)$$

where g is the determinant of (g_{ab}) . Note that for a unimodular, in particular semisimple, group $C_{ab}^b = 0$, and the second term on the right hand side of (81) vanishes. The corresponding Bianchi models are known as Class A models.

As seen from Eq. (80), $\hat{D}_1$ acts by multiplication by a time-dependent function. Therefore, choosing $q = i\sqrt{g}$ reduces Eq. (14) to the eigenvalue equation:

$$(\hat{D}_2 - E_n^2) \Phi_n = -(\Delta_t + E_n^2 - \mu^2) \Phi_n = 0, \quad (82)$$

for the operator $\hat{D}_2$ which being essentially the Laplacian over Σ_t , is self-adjoint. This guarantees the orthogonality of Φ_n and the reality of E_n^2 . Furthermore, since q is imaginary, the Hamiltonian $H^{(q)}$ with the choice of (11) for the inner product is also self-adjoint.

The analysis of the Φ_n is equivalent to the study of the eigenvectors of the Laplacian over a three-dimensional group manifold Σ_t . The general problem is the subject of the investigation in spectral geometry which is beyond the scope of the present article. However, let us recall some well-known facts about spectral properties of the Laplacian Δ for an arbitrary finite-dimensional Riemannian manifold Σ without boundary.

The following results are valid for the case where Σ is compact or the eigenfunctions are required to have a compact support⁶ [35]:

⁶This is equivalent with the case where Σ has a boundary $\partial\Sigma$, over which the eigenfunctions vanish.

1. The spectrum of Δ is an infinite discrete subset of non-negative real numbers.
2. The eigenvalues are either non-degenerate or finitely degenerate.
3. There is an orthonormal set of eigenfunctions which form a basis for $L^2(\Sigma)$.
4. If Σ is compact, then the first eigenvalue is zero which is non-degenerate with the eigenspace given by the set of constant functions, i.e., $\mathbb{C}$. If Σ is not compact but the eigenfunctions are required to have a compact support, then the first eigenvalue is positive.

In view of Eq. (82) and the results concerning the spectral properties of the Laplacian Δ_t , one has some restrictions on the energy eigenvalues E_n . The non-negativity of the eigenvalues $\mu^2 - E_n^2$ of Δ_t is a necessary condition for the validity of the first quantized theory. A violation of this condition is equivalent to the Klein paradox [24]. In fact, similar conditions exist for both rotating magnetic field and rotating cosmic string problems. In these cases, it indicates that the variables k appearing in the analysis of these systems must be imaginary.

Another piece of useful information about the spatially homogeneous cosmological models is that Σ_t is a group manifold, i.e., topologically it is identical to a three-dimensional Lie group G . Therefore, one can use the canonical (left and right) invariant metric (δ_{ab}) on G , to define the corresponding Laplacian Δ_0 and use an orthonormal eigenbasis f_n of Δ_0 to span $L^2(\Sigma_t) = \mathcal{H}_t$, i.e., express Φ_n as a linear combination of f_n . Perhaps more importantly, using the fact that (up to a multiple of $i = \sqrt{-1}$) the invariant vector fields $\hat{X}_a$ yield a representation of the generators L_a of G , with $L^2(\Sigma_t)$ being the representation space, one can view the Laplacian Δ_t as (a representation of) an element of the enveloping algebra of the Lie algebra of G . Therefore, Δ_t commutes with any Casimir operator $\mathcal{C}_\lambda$ and consequently shares a set of simultaneous eigenvectors with $\mathcal{C}_\lambda$. This in turn suggests one to specialize to particular subrepresentations with definite $\mathcal{C}_\lambda$. In particular for compact groups, this leads to a reduction of the problem to a collection of finite-dimensional ones.⁷

In the remainder of this article I shall try to employ these considerations to investigate some specific models.

⁷Here I mean a finite-dimensional Hilbert space.

6.1 Bianchi Type I

In this case G is Abelian, therefore X_a are themselves Casimir operators and the eigenfunctions of Δ_t , i.e., Φ_n , are independent of t . Hence the Berry connection one-form (31) vanishes identically and the geometric phase is trivial.⁸

6.2 Bianchi Type IX

In this case $G = SU(2)$. The total angular momentum operator $\hat{J}^2 = \sum_a \hat{J}_a^2$ is a Casimir operator. Therefore, I shall consider the subspaces $\mathcal{H}_j$ of $\mathcal{H}_t = L^2(S_t^3)$ of definite angular momentum j . The left-invariant vector fields $\hat{X}_a$ are given by $\hat{X}_a = i\hat{J}_a$, in terms of which Eq. (77), with $C_{ab}^c = \epsilon_{abc}$, is written in the familiar form:

$$[\hat{J}_a, \hat{J}_b] = i\epsilon_{abc}\hat{J}_c, \quad (83)$$

with ϵ_{abc} denoting the totally antisymmetric Levi Civita symbol and $\epsilon_{123} = 1$.

Eq. (82) takes the form:

$$(\hat{H}' + k_n^2)\Phi_n = 0, \quad (84)$$

where $\hat{H}'$ is an induced Hamiltonian defined by

$$\hat{H}' := g^{ab}(t)\hat{J}_a\hat{J}_b, \quad (85)$$

and $k_n^2 := E_n^2 - \mu^2$. Therefore, the problem of the computation of the geometric phase is identical with that of the non-relativistic quantum mechanical system whose Hamiltonian is given by (85). In particular, for the mixmaster spacetime, i.e., for g_{ab} diagonal, the problem is identical with the quantum mechanical problem of a non-relativistic asymmetric rotor, [36].

Another well-known non-relativistic quantum mechanical effect which is described by a Hamiltonian of the form (85) is the quadratic interaction of a spin with a variable electric field (E_a). The interaction potential is the Stark Hamiltonian: $\hat{H}_S = \epsilon(\sum_a E_a \hat{J}_a)^2$. The phenomenon of the geometric phase for the Stark Hamiltonian for spin $j = 3/2$, which involves Kramers degeneracy [26], was first considered by Mead [37]. Subsequently, Avron, et al [38, 39] conducted a thorough investigation of the traceless quadrupole Hamiltonians of the form (85).

⁸By a trivial geometric phase, I mean a zero geometrical phase angle.

The condition on the trace of the Hamiltonian is physically irrelevant, since the addition of any multiple of the identity operator to the Hamiltonian does not have any physical consequences. In general, one can express the Hamiltonian (85) in the form $\hat{H}' = \hat{\hat{H}}' + \hat{H}'_0$, where $\hat{\hat{H}}' := Tr(g^{ab})\hat{J}^2/3$,

$$\hat{H}'_0[R] := \sum_{A=1}^5 R^A \hat{e}_A, \quad (86)$$

is the traceless part of the Hamiltonian, and

$$\begin{aligned} \hat{e}_1 &:= J_3^2 - \frac{1}{3} \hat{J}^2, & \hat{e}_2 &:= \frac{1}{\sqrt{3}} \{\hat{J}_1, \hat{J}_2\}, \\ \hat{e}_3 &:= \frac{1}{\sqrt{3}} \{\hat{J}_2, \hat{J}_3\}, & \hat{e}_4 &:= \frac{1}{\sqrt{3}} (\hat{J}_1^2 - \hat{J}_2^2), \\ \hat{e}_5 &:= \frac{1}{\sqrt{3}} \{\hat{J}_1, \hat{J}_2\}, \end{aligned} \quad (87)$$

$$\begin{aligned} R^1 &:= g^{33} - \frac{1}{2} (g^{11} + g^{22}), & R^2 &:= \sqrt{3} g^{13}, \\ R^3 &:= \sqrt{3} g^{23}, & R^4 &:= \frac{\sqrt{3}}{2} (g^{11} - g^{22}), \\ R^5 &:= \sqrt{3} g^{12}. \end{aligned} \quad (88)$$

As shown in Refs. [38, 39] the space $\mathcal{M}'$ of all traceless Hamiltonians of the form (86) is a five-dimensional real vector space. The traceless operators $\hat{e}_A$ form an orthonormal⁹ basis for $\mathcal{M}'$. Removing the point ($R^A = 0$) from $\mathcal{M}'$, to avoid the collapse of all eigenvalues, one has therefore the space $\mathbb{R}^5 - \{0\}$ as the parameter space. The situation is analogous to Berry's original example of a magnetic dipole in a changing magnetic field, [1]. Again, a rescaling of the Hamiltonian by a non-zero function of R^A does not change the geometric phase. Thus the relevant parameter space is $\mathcal{M} = S^4$. Incidentally, the point corresponding to $0 \in \mathbb{R}^5$ which is to be excluded, corresponds to the class of Friedmann-Robertson-Walker models.

Clearly a curve $C : [0, T] \rightarrow S^4$ may be defined by the action of the group $SO(5)$ which acts transitively on S^4 . Therefore the time-dependence of the Hamiltonian may be realized by an action of the group $SO(5)$ on a fixed Hamiltonian. As it is discussed in Ref. [39], it is the unitary representations $\mathcal{U}$ of the double cover $Spin(5) = Sp(2)$ of $SO(5)$ (alternatively the projective representations of $SO(5)$) which define the time-dependent Hamiltonian:

$$\hat{H}'_0[R(t)] = \mathcal{U}[g(t)] \hat{H}'_0[R(0)] \mathcal{U}[g(t)]^\dagger. \quad (89)$$

⁹Orthonormality is defined by the inner product $\langle \hat{A}, \hat{B} \rangle := 3 Tr(\hat{A}\hat{B})/2$.

Here, $g : [0, T] \rightarrow Sp(2)$, is defined by $R(t) =: \pi[g(t)]R(0)$, where $\pi : Sp(2) \rightarrow SO(5)$ is the canonical two-to-one covering projection. The emergence of the group $Sp(2)$ is an indication of the existence of a quaternionic description of the system, [39].

Let us next examine the situation for irreducible representations j of $SU(2)$. As I previously described, $\hat{J}^2$ commutes with the Hamiltonian. Hence the Hamiltonian is block-diagonal in the basis with definite total angular momentum j . For each j , the representation space $\mathcal{H}_j$ is $2j + 1$ dimensional. Therefore the restriction of the Hamiltonian $\hat{H}'_0$ to $\mathcal{H}_j$ and $\mathcal{U}[g(t)]$ are $(2j + 1) \times (2j + 1)$ matrices.

Let $\{\phi_{j_n}^I\}$ be a complete set of orthonormal eigenvectors of the initial Hamiltonian $\hat{H}'_0[R(0)]$, where I is a degeneracy label. Then the eigenvectors of $\hat{H}'_0[R]$ are of the form:

$$\Phi_{j_n}^I[R] = \mathcal{U}[g] \phi_{j_n}^I, \quad (90)$$

and the non-Abelian connection one-form (24) is given by

$$\mathcal{A}_{j_n}^{IJ} = i \langle \phi_{j_n}^I | \mathcal{U}^\dagger d\mathcal{U} | \phi_{j_n}^J \rangle, \quad (91)$$

where d stands for the exterior derivative operator with respect to the parameters of the system.

For the integer j , bosonic systems, it is known that the quadratic Hamiltonians of the form (86), describe time-reversal-invariant systems. In this case it can be shown that the curvature two-form associated with the Abelian Berry connection one-form (31) vanishes identically [39]. The connection one-form is exact (gauge potential is pure gauge) and a nontrivial geometric phase can only be topological, namely it may still exist provided that the first homology group of the parameter space is nontrivial. For the problem under investigation $\mathcal{M} = S^4$, and the first homology group is trivial. Hence, in general, Abelian geometric phases are trivial. The same conclusion cannot however be reached for the non-Abelian (matrix-valued) geometric phases.

In the remainder of this section, I shall examine the situation for some small values of j :

- 1) $j = 0$: The corresponding Hilbert subspace is one-dimensional. Therefore, there is no nontrivial geometric phases.
- 2) $j = 1/2$: In this case, $\hat{J}_a = \sigma_a/2$, where σ_a are Pauli matrices. Using the well-known anticommutation (Clifford algebra) relations $\{\sigma_a, \sigma_b\} = 2\delta_{ab}$, one can easily show that in

this case

$$\hat{H}' = \frac{1}{2} \sum_a g^{aa}(t) \hat{I} , \quad (92)$$

where $\hat{I}$ is the identity matrix. Therefore, the eigenvectors Φ_n are constant (g_{ab} -independent), the connection one-form (91) vanishes and the geometric phase angle is again zero.

- 3) $j = 1$: In this case the Hilbert subspace is three-dimensional. The Abelian geometrical phases are trivial. The nontrivial matrix-valued geometrical phases may be present, provided that the Hamiltonian has a degenerate eigenvalue. Using the ordinary $j = 1$ matrix representations of $\hat{J}_a$, one can easily express $\hat{H}'_0$ as a 3×3 matrix. It can then be checked that in the generic case the eigenvalues of $\hat{H}'_0$ are not degenerate.¹⁰ However, there are cases for which a degenerate eigenvalue is present. A simple example is the Taub metric, $(g_{ab}) = \text{diag}(g_{11}, g_{22}, g_{33})$, which is a particular example of the mixmaster metric, [34]. For the general mixmaster metric, the eigenvalue problem can be easily solved. The eigenvalues of the total Hamiltonian $\hat{H}'$ of Eq. (85) are

$$\frac{1}{g_{11}} + \frac{1}{g_{22}} , \quad \frac{1}{g_{11}} + \frac{1}{g_{33}} , \quad \frac{1}{g_{22}} + \frac{1}{g_{33}} .$$

Therefore the degenerate case corresponds to the coincidence of at least two of g_{aa} 's, i.e., a Taub metric. However, even in the general mixmaster case, one can find a constant (g_{aa} -independent) basis which diagonalizes the Hamiltonian. Hence the non-Abelian connection one-form vanishes and the geometric phase is again trivial. This is not however the case for general metrics. In Appendix B, it is shown that without actually solving the general eigenvalue problem for the general Hamiltonian, one can find the conditions on the metric which render at least one of the eigenvalues of the Hamiltonian degenerate. Here, I summarize the results. Using the well-known matrix representations of the angular momentum operators $\hat{J}_a$ in the $j = 1$ representation [40] one can write the Hamiltonian (85) in the form:

$$\hat{H}' = \begin{pmatrix} t + 2z & \xi^* & \zeta^* \\ \xi & 2(t + z) & -\xi^* \\ \zeta & -\xi & t + 2z \end{pmatrix} , \quad (93)$$

¹⁰This is true for all integer j , [39].

where

$$\begin{aligned} t &:= \frac{1}{2} (g^{11} + g^{22}) - g^{33} , & z &:= g^{33} , \\ \xi &:= \frac{1}{\sqrt{2}} (g^{13} + ig^{23}) , & \zeta &:= \frac{1}{2} (g^{11} - g^{22}) + ig^{12} . \end{aligned}$$

Then it can be shown (App. B), that the necessary and sufficient conditions for $\hat{H}'$ to have a degenerate eigenvalue are

I. for $\zeta = 0$: $\xi = 0$, in which case, $\hat{H}'$ as given by Eq. (93) is already diagonal. The degenerate and non-degenerate eigenvalues are $t + 2z$ and $2(t + z)$, respectively. In terms of the components of the metric, these conditions can be written as: $g_{11} = g_{22}$ and $g_{ab} = 0$ if $a \neq b$. This is a Taub metric which as discussed above does not lead to a nontrivial geometric phase.

II. for $\zeta \neq 0$:

$$\zeta = \mathcal{Z} e^{2i\theta} , \quad t = \mathcal{Z} - |\xi|^2/\mathcal{Z} , \quad (94)$$

where $\exp[i\theta] := \xi/|\xi|$ and $\mathcal{Z} \in \mathbb{R} - \{0\}$. In this case the degenerate and non-degenerate eigenvalues are $2(\mathcal{Z} + z) - |\xi|^2/\mathcal{Z}$ and $2(z - |\xi|^2/\mathcal{Z})$, respectively.

For the latter case, an orthonormal set of eigenvectors is given by:

$$v_1 = \frac{1}{\sqrt{2}} \begin{pmatrix} e^{-2i\theta} \\ 0 \\ 1 \end{pmatrix} , \quad v_2 = \frac{1}{\sqrt{1+2\mathcal{X}^2}} \begin{pmatrix} \mathcal{X}e^{-i\theta} \\ 1 \\ -\mathcal{X}e^{i\theta} \end{pmatrix} , \quad v_3 = \frac{1}{\sqrt{2(1+2\mathcal{X}^2)}} \begin{pmatrix} -e^{-2i\theta} \\ 2\mathcal{X}e^{-i\theta} \\ 1 \end{pmatrix} , \quad (95)$$

where $\mathcal{X} := |\xi|/(2\mathcal{Z})$. In view of the general argument valid for all non-degenerate eigenvalues, the geometric phase associated with v_3 is trivial. This can be directly checked by substituting v_3 in the formula (31) for the Berry connection one-form. This leads, after some algebra, to the surprisingly simple result $\mathcal{A}_{33} := i\langle v_3 | dv_3 \rangle = d\theta$. Therefore, $\mathcal{A}_{33}$ is exact as expected, and the corresponding geometric phase angle vanishes. Similarly one can compute the matrix elements $\mathcal{A}_{rs} := i\langle v_r | dv_s \rangle$, $r, s = 1, 2$, of the non-Abelian connection one-form. The result is:

$$\mathcal{A} = \begin{pmatrix} 1 & \mathcal{F} \\ \mathcal{F}^* & 0 \end{pmatrix} \omega , \quad (96)$$

$$\begin{aligned}\mathcal{F} &:= \frac{2\mathcal{X} e^{i\theta}}{\sqrt{2(1+2\mathcal{X}^2)}} = \frac{2\epsilon\xi}{\sqrt{2+|\frac{\xi}{\zeta}|^2}} = \frac{\epsilon(g^{13}+ig^{23})}{\sqrt{1+\frac{(g^{13})^2+(g^{23})^2}{(g^{11}-g^{22})^2+(2g^{12})^2}}}, \\ \omega &:= d\theta = \frac{g^{13}dg^{23}-g^{23}dg^{13}}{(g^{13})^2+(g^{23})^2},\end{aligned}$$

where $\epsilon := \mathcal{Z}/|\zeta| = \pm 1$. As seen from Eq. (96), $\mathcal{A}$ is a $u(2)$ -valued one-form, which vanishes if g^{23}/g^{13} is kept constant during the evolution of the universe.

It is also worth mentioning that the requirement of the existence of degeneracy is equivalent to restricting the parameters of the system to a two-dimensional subset of S^4 . Thus, the corresponding spectral bundle [27, 28] is a $U(2)$ vector bundle over a two-dimensional parameter space $\tilde{\mathcal{M}}$. The manifold structure of $\tilde{\mathcal{M}}$ is determined by Eqs. (94). In terms of the parameters R^A of (88), these equations are expressed by

$$R^5 = f_1 R^4, \quad R^1 = f_2 R^4 + \frac{f_3}{R^4}, \quad (97)$$

where

$$\begin{aligned}f_1 &:= \frac{2R^2 R^3}{(R^2)^2 - (R^3)^2}, & f_2 &:= \pm \frac{(R^2)^2 + (R^3)^2}{\sqrt{3}[(R^2)^2 - (R^3)^2]}, \\ f_3 &:= \mp \frac{(R^2)^2 - (R^3)^2}{2\sqrt{3}}, & f_4 &:= (R^2)^2 + (R^3)^2.\end{aligned}$$

Here f_4 is also introduced for future use. In addition to (97), one also has the condition $(R^A) \in S^4$. If S^4 is identified with the round sphere, this condition takes the form $\sum_A (R^A)^2 = 1$. Substituting (97) in this equation, one finds

$$(1 + f_2^2 + f_3^2)(R^4)^4 - (1 - f_4 - 2f_2 f_3)(R^4)^2 + f_3^2 = 0. \quad (98)$$

Eq. (98) may be easily solved for R^4 . This yields:

$$R^4 = \pm \frac{3[(R^2)^2 - (R^3)^2]^2}{8[(R^2)^2 + (R^3)^2]^2} \left[1 - \frac{2}{3}[(R^2)^2 + (R^3)^2] \pm \sqrt{1 - \frac{4}{3}[(R^2)^2 + (R^3)^2]} \right]. \quad (99)$$

Note that the parameters R^A are related to the components of the inverse of the three-metric through Eqs. (88). Thus the parameter space $\tilde{\mathcal{M}}$ is really a submanifold of the corresponding minisuperspace. Fig. 1 shows a three-dimensional plot of R^4 as a function of R^2 and R^3 , i.e., a plot of the parameter space $\tilde{\mathcal{M}}$ as embedded in $\mathbb{R}^3$. Note that

$R^2 = \pm R^3$ renders f_1 and f_2 singular. The corresponding points which are depicted as the curves along which the figure becomes non-differentiable must be handled with care. The smooth part of $\tilde{\mathcal{M}}$ consists of eight connected components, each of which is diffeomorphic to an open disk (alternatively $\mathbb{R}^2$).

- 4) $j = 3/2$: This case has been studied in Refs. [38, 39] in detail. Therefore I suffice to note that it involves nontrivial geometric phases.

7 Conclusion

In this article I showed that the two-component formalism could be consistently used to investigate the geometric phases associated with charged Klein-Gordon fields. This formalism provides a precise definition of the adiabatic approximation and allows Berry's derivation of the adiabatic geometrical phase to be applied to the relativistic Klein-Gordon fields. In particular, I showed that the computation of the adiabatic geometric phase did not involve the explicit construction of an inner product on the space of the initial conditions, or alternatively the space of solutions of the Klein-Gordon equation. It only required the inner product structure of the Hilbert space $L^2(\Sigma_t)$.

In non-relativistic quantum mechanics, the necessary and sufficient condition for the validity of the adiabatic approximation, $\psi \approx e^{i\alpha}|n\rangle$, is $\langle m|\dot{n}\rangle \approx 0$ for $m \neq n$, [26, 30], where $|n\rangle$ are instantaneous eigenvectors of the Hamiltonian. If the Hamiltonian is not self-adjoint then the eigenvectors may not be orthogonal. In this case this condition is generalized to $\langle n|n\rangle\langle m|\dot{n}\rangle - \langle m|n\rangle\langle n|\dot{n}\rangle \approx 0$. A direct generalization of the ansatz $\psi \approx e^{i\alpha}|n\rangle$ within the two-component formulation of the Klein-Gordon equation leads to an additional condition on the energy eigenvalues, namely $\frac{d(qE_n)}{dt} \approx 0$. If this condition is satisfied then the evolution is said to be ultra-adiabatic. If this condition fails to be fulfilled but the adiabaticity condition (35) is satisfied, then the evolution is said to be adiabatic. The expressions for the geometric phase for the ultra-adiabatic and adiabatic evolutions are identical. The only difference is in the dynamical part of the phase.

I employed the general results of the two-component formulation to study adiabatic geo-

metric phases induced by a rotating magnetic field and a rotating cosmic string. The results were in complete agreement with those of the previous investigations [6, 17], but the analysis was considerably simpler.

I also investigated the geometric phases induced by spatially homogeneous cosmological backgrounds. In this case the freedom in the choice of q turned out to simplify the analysis. I showed that the adiabatic geometric phase angles always vanished for Bianchi type I models, whereas non-Abelian adiabatic phases could occur for the Bianchi type IX models. Particularly, interesting was the relationship between the corresponding induced Hamiltonians in the Bianchi type IX models and the quadrupole Hamiltonians of the molecular and nuclear physics. I also showed that even for the integer spin representations nontrivial geometric phases could exist. This should also be of interest for the molecular physicists and chemists who have apparently investigated only the fermionic systems (half-integer spin representations.)

In the context of general relativity where the Poincaré invariance is replaced by the diffeomorphism invariance, one can use the time-reparameterization symmetry of the background gravitational field and the geometric phase to absorb the magnitude $|q|$ of the decomposition parameter q into the definition of the lapse function $N = (-g^{00})^{-1/2}$. In this way only a $U(1)$ subgroup of the corresponding $GL(1, \mathbb{C})$ symmetry group survives. The additional requirement of the self-adjointness of the Hamiltonian with respect to the Klein-Gordon inner product, can then be used to choose q to be imaginary, i.e., to set $q = i$ or $q = -i$. Note however that the Klein-Gordon inner product does not play any distinguished role in the computation of the geometric phase. Hence the choice $q = i$ or $q = -i$ is not a necessary condition. It should also be clear that the $GL(1, \mathbb{C})$ or $U(1)$ symmetry associated with the freedom of choice of the decomposition has no physical basis or consequences. It is merely a mathematical feature of the two-component formalism which can occasionally be used to simplify the calculations.

Finally, I wish to emphasize that the use of the two-component formulation in the study of the geometric phases associated with scalar fields is more advantageous than the more conventional approaches which are based on a decomposition of the space of solutions into positive and negative frequency subspaces and the construction of a positive definite inner product, e.g., those used in Refs. [6, 17]. This has two reasons. Firstly, the conventional methods have

apparently missed the fact that one does not need to construct an inner product on the space of Klein-Gordon fields to be able to calculate the adiabatic geometric phase. Hence, a major part of these analyses is concerned with the construction of such an inner product. Secondly, these approaches can only be applied to the stationary spacetimes where such an inner product can be constructed. The application of the two-component formulation for the Bianchi models manifestly shows that this method can also be employed even if the background spacetime is not stationary. One must however realize that similarly to the conventional methods the present analysis is only valid within the framework of the adiabatic approximation. Although, the (approximate) stationarity of the background metric (for $A = 0 = V$) is a sufficient condition for the validity of the adiabatic approximation, it is not necessary. This can be easily seen by noting that for example in the case of Bianchi IX model, for spin $j = 1/2$ states, one has $\dot{\Phi}_n = 0$, so $\langle \Phi_m | \dot{\Phi}_n \rangle = 0$. Therefore, although the spacetime is not stationary, the adiabatic approximation yields the exact solution of the field equation. This shows that in general for arbitrary non-stationary spacetimes, there may exist adiabatically evolving states to which the above analysis applies. This is in contrast with the traditional methods of the computation of the adiabatic geometric phases for scalar fields which involve construction of a Hilbert space structure on the space of solutions of the Klein-Gordon equation.

Acknowledgements

I would like to thank Bahman Darian for many fruitful discussions and helping me with the computer graphics, Teoman Turgut for bringing to my attention one of the references, and Kamran Saririan for mailing me copies of a couple of the references.

Appendix A

In this Appendix I show that for the problem of a rotating cosmic string, there exist localized energy eigenvectors.

Let $f = f(\rho, \varphi)$ be an arbitrary localized smooth function, i.e., it has a compact support,

and write

$$f(\rho, \varphi) = \sum_m \int dk \tilde{f}_m(k) e^{im\varphi} J_\nu(k\rho) , \quad (100)$$

where $\nu = m/\alpha$. Using the orthogonality properties of the Bessel functions and the ‘plane waves’, Eq. (100) may be easily inverted to yield $\tilde{f}_m(k)$ which in turn define a localized field with definite energy eigenvalue E_n according to:

$$\Phi_n = e^{-4iE_n j\varphi} \sum_m \int dk \tilde{f}_m(k) e^{im\varphi} e^{i(\sqrt{E_n^2 - \mu^2 - k^2}z)} J_\nu(k\rho) .$$

By construction Φ_n is localized in the ρ and φ -directions. The localization in z -direction is irrelevant to the arguments used in section 5. Furthermore, note that there may still be degenerate degrees of freedom left in this construction.

Appendix B

In this Appendix I show how one can obtain the conditions under which the Hamiltonian (93) has degenerate eigenvalues without actually solving the eigenvalue problem in the general case.

The analysis can be slightly simplified if one writes the Hamiltonian (93) in the form:

$$\begin{aligned} \hat{H}' &= (t + 2z)\hat{I} + \hat{\tilde{H}} , \\ \hat{\tilde{H}} &:= \begin{pmatrix} 0 & \xi^* & \zeta^* \\ \xi & t & -\xi^* \\ \zeta & -\xi & 0 \end{pmatrix} , \end{aligned} \quad (101)$$

where $\hat{I}$ is the 3×3 identity matrix. Clearly, the eigenvalue problems for $\hat{H}'$ and $\hat{\tilde{H}}$ are equivalent. Computing the characteristic polynomial for $\hat{\tilde{H}}$, i.e., $P(\lambda) := \det(\hat{\tilde{H}} - \lambda\hat{I})$, one finds:

$$P(\lambda) = -\lambda^3 + t\lambda^2 + (|\zeta|^2 + 2|\xi|^2)\lambda - (t|\zeta|^2 + \zeta\xi^{*2} + \zeta^*\xi^2) . \quad (102)$$

If one of the eigenvalues (roots of $P(\lambda)$) is degenerate, then

$$P(\lambda) = -(\lambda - l_1)(\lambda - l_2)^2 . \quad (103)$$

Comparing Eqs. (102) and (103), one finds

$$t = l_1 + 2l_2 , \quad l_2^2 + 2l_1l_2 = -(|\zeta|^2 + 2|\xi|^2) , \quad l_1l_2^2 = -(t|\zeta|^2 + \zeta\xi^{*2} + \zeta^*\xi^2) . \quad (104)$$

Furthermore since l_2 is at least doubly degenerate, the rows of the matrix:

$$\hat{H} - l_2 \hat{I} = \begin{pmatrix} -l_2 & \xi^* & \zeta^* \\ \xi & t - l_2 & -\xi^* \\ \zeta & -\xi & -l_2 \end{pmatrix}, \quad (105)$$

must be mutually linearly dependent. In other words the cofactors of all the matrix elements must vanish. Enforcing this condition for the matrix elements and using Eqs. (104), one finally finds that either $\xi = \zeta = l_2 = 0$ and $l_1 = t$, or

$$\begin{aligned} \zeta &= \mathcal{Z} e^{2i\theta}, & t &= \mathcal{Z} - \frac{|\xi|^2}{\mathcal{Z}}, \\ l_2 &= \mathcal{Z}, & l_1 &= -(\mathcal{Z} + \frac{|\xi|^2}{\mathcal{Z}}), \end{aligned}$$

where $\exp[i\theta] := \xi/|\xi|$ and $\mathcal{Z} \in \mathbb{R} - \{0\}$.


References

- [1] M. V. Berry, Proc. Roy. Soc. London **A392**, 45 (1984).
- [2] F. Wilczek and A. Zee, Phys. rev. Lett. **52**, 2111 (1984).
- [3] Y. Aharonov and J. Anandan, Phys. Rev. Lett. **58**, 1593 (1987); J. Anandan and Y. Aharonov, Phys. Rev. **D38**, 1863 (1988).
- [4] J. C. Garrison and R. Y. Chiao, Phys. Rev. Lett. **60**, 165 (1988).
- [5] J. Anandan, Phys. Rev. Lett. **60**, 2555 (1988).
- [6] J. Anandan and P. O. Mazur, Phys. Lett. **A173**, 116 (1993).
- [7] Ph. Droz-Vincent, Lett. Math. Phys. **36**, 277 (1996).
- [8] R. Brout and G. Venturi, Phys. Rev. **D39**, 2436 (1989).
- [9] T. Banks, Nucl. Phys. **B249**, 332 (1985).
- [10] R. Brout, Found. Phys. **17**, 603 (1987).
- [11] J. Moody, A. Shapere, and F. Wilczek, Phys. Rev. Lett. **56**, 893 (1986).

- [12] G. Venturi, “Quantum Gravity and the Berry Phase,” in *Differential Geometric Methods in Theoretical Physics*, Eds. L.-L. Chau and W. Nahm, Plenum, New York, 1990.
- [13] G. Venturi, *Class. Quantum Grav.* **7**, 1075 (1990).
- [14] R. Casadio and G. Venturi, *Class. Quantum Grav.* **12**, 1267 (1995).
- [15] D. P. Datta, *Phys. Rev.* **D 48**, 5746 (1993); *Gen. Rel. Grav.* **27**, 341 (1995).
- [16] Y. Q. Cai and G. Papini, *Mod. Phys. Lett.* **A4**, 1143 (1989); *Class. Quantum Grav.* **7**, 269 (1990).
- [17] A. Corichi and M. Pierri, *Phys. Rev.* **D51**, 5870 (1995).
- [18] P. de Sousa Gerbert and J. Jackiw, *Commun. Math. Phys.* **124**, 229 (1989).
- [19] B. S. DeWitt, *Phys. Rep.* **C19**, 295 (1975).
- [20] S. A. Fulling, *Aspects of Quantum Field Theory in Curved Space-Time*, Cambridge University Press, Cambridge, 1989.
- [21] R. M. Wald, *Quantum Field Theory in Curved Spacetime and Black Hole Thermodynamics*, The University of Chicago Press, Chicago, 1994.
- [22] C. W. Misner, K. S. Thorne, and J. A. Wheeler, *Gravitation*, Freeman, New York, 1973.
- [23] H. Feshbach and F. Villars, *Rev. Mod. Phys.* **30**, 24 (1958).
- [24] B. R. Holstein, *Topics in Advanced Quantum Mechanics*, Addison-Wesley, Redwood City, California, 1992.
- [25] R. M. Wald, *General Relativity*, The University of Chicago Press, Chicago, 1984.
- [26] A. Messiah, *Quantum Mechanics*, Vol. 2, North-Holland, Amsterdam, 1962; L. I. Schiff, *Quantum Mechanics*, McGraw-Hill, New York, 1955.
- [27] B. Simon, *Phys. Rev. Lett.* **51**, 2167, (1983).

- [28] A. Bohm and A. Mostafazadeh, J. Math. Phys. **35**, 1463 (1994).
- [29] J. Samuel and R. Bhandari, Phys. Rev. Lett. **60**, 2339 (1988).
- [30] A. Mostafazadeh, Phys. Rev. **A 55**, 1653 (1997).
- [31] A. Bohm, *Quantum Mechanics: Foundations and Applications*, third edition, Springer-Verlag, Berlin, 1993.
- [32] Y. Aharonov and D. Bohm, Phys. Rev. **115**, 485 (1959).
- [33] A. Ashtekar and A. Mangon, Proc. R. Soc. London **A346**, 375 (1975).
- [34] M. P. Ryan, Jr. and L. C. Shepley, *Homogeneous Relativistic Cosmologies*, Princeton University Press, Princeton, 1975.
- [35] S. Gallot, D. Hulin, and J. Lafontaine, *Riemannian Geometry*, Springer-Verlag, Berlin, 1987.
- [36] B. L. Hu, Phys. Rev. **D8**, 2377 (1973).
- [37] C. A. Mead, Phys. Rev. Lett. **59**, 161 (1987).
- [38] J. E. Avron, L. Sadun, J. Segret, and B. Simon, Phys. Rev. Lett. **61**, 1329 (1988).
- [39] J. E. Avron, L. Sadun, J. Segret, and B. Simon, Commun. Math. Phys. **124**, 595 (1989).
- [40] L. I. Schiff, *Quantum Mechanics*, McGraw-Hill, New York, 1955.

Figure 1: This is a plot of $R^4 = R^4(R^2, R^3)$. The horizontal plane is the R^2 - R^3 -plane and the vertical axis is the R^4 -axis. The parameter space $\tilde{\mathcal{M}}$ is obtained by removing the intersection of this figure with the planes defined by: $R^4 = 0$, $R^2 = R^3$ and $R^2 = -R^3$. The intersection involves the curves along which the figure becomes non-differentiable.



This figure "fig1-1.png" is available in "png" format from:

<http://arXiv.org/ps/hep-th/9608051v3>